

Comparison of Receptor Profiles Between Primary Tumors and Isolated Thoracic Metastases in Breast Cancer

Comparação dos Perfis de Recetores entre os Tumores Primários e as Metástases Torácicas Isoladas no Cancro da Mama

Aybiyçe Elif Silpağar Güner ¹, Eyüp Halit Yardımcı ², Mehmet Yıldırım ³

1 - Department of Thoracic Surgery, Sancaktepe Şehit Prof. Dr. İlhan Varank Training and Research Hospital, İstanbul, Türkiye .

2 - Department of Thoracic Surgery, Marmara University, School of Medicine, İstanbul, Türkiye

3 - Department of Thoracic Surgery, Dr. Siyami Ersek Training and Research Hospital, İstanbul, Türkiye

<https://doi.org/10.82582/thorac.95>

Autor Correspondente/Corresponding Author:

Dr. Aybiyçe Elif Silpağar Güner

<https://orcid.org/0000-0003-2149-9077>

Email: aybiycesilpagar@gmail.com

Declaração de Contribuição/Contributorship Statement:

Todos os autores contribuíram para este trabalho, aprovaram a versão final do manuscrito para publicação e concordam em assumir a responsabilidade por todos os aspetos do trabalho, garantindo a exatidão e a integridade dos dados apresentados.

All authors contributed to this work, approved the final version of the manuscript for publication, and agree to be accountable for all aspects of the work, ensuring the accuracy and integrity of the data presented.

Recebido/Received: May 11th, 2026

Aceite/Accepted: May 14th, 2026

Publicado online/Published online: August 14th, 2026

Publicado / Published: August 18th, 2026

© 2026 THORAC. This is an open-access article under the CC BY-NC 4.0. Re-use permitted under CC BY-NC 4.0. No commercial re-use.

© 2026 THORAC. Este é um artigo de acesso aberto sob a licença CC BY-NC 4.0. Reutilização permitida de acordo com CC BY-NC 4.0.

Nenhuma reutilização comercial.

ABSTRACT

Introduction

Receptor status may change during metastatic progression, potentially altering therapeutic decisions and outcomes. This study aimed to determine ER, PR, and HER2 changes in thoracic metastases (pleura, lung, lymph node, pericardium, chest wall) of breast cancer and to evaluate their impact on survival.

Methods

Data of 46 women who underwent an interventional procedure for thoracic metastasis of breast cancer and had available ER/PR/HER2 assessment in both primary and corresponding metastatic tissues were retrospectively analyzed. Post-metastasis survival was calculated from the date of histopathological confirmation of metastatic disease. The associations of primary breast cancer receptor status, metastatic receptor status, and receptor conversion at metastasis with survival outcomes were investigated.

Results

The pleura was the most common metastatic site (54,3%). In primary tumors, ER, PR, and HER2 positivity rates were 87%, 63%, and 21%, respectively, whereas in metastatic tissues they were 69,6%, 45,7%, and 25,6%. ER loss was observed in 17,4% of cases, with no gain detected. PR loss occurred in 23,9% and gain in 6,5%, while HER2 loss and gain were observed in 7,0% and 11,6% of cases, respectively. Univariate analysis identified pleural metastasis (HR=3,442, $p=0,017$), ER negativity in metastatic tissue (HR=3,306, $p=0,008$), PR negativity in metastatic tissue (HR=2,793, $p=0,037$), and ER loss (HR=3,095, $p=0,022$) as adverse prognostic factors. In multivariate analysis, pleural metastasis (HR=4,424, $p=0,009$) and ER loss (HR=3,669, $p=0,010$) remained independent predictors of poor survival. No significant association was found between HER2 status and survival.

Conclusion

ER loss and pleural metastasis are independent adverse prognostic factors in thoracic metastases of breast cancer. These findings related to tumor biology, particularly receptor status changes between primary and metastatic tumors, may help guide personalized therapeutic approaches.

terapêuticas e os resultados clínicos. Este estudo teve como objetivo determinar as alterações dos recetores de estrogénio (ER), progesterona (PR) e HER2 nas metástases torácicas (pleura, pulmão, gânglios linfáticos, pericárdio e parede torácica) do cancro da mama, bem como avaliar o seu impacto na sobrevivência.

Métodos

Foram analisados retrospectivamente os dados de 46 mulheres submetidas a um procedimento de intervenção para metástases torácicas de cancro da mama, nas quais estavam disponíveis as determinações de ER, PR e HER2 tanto no tumor primário como no tecido metastático correspondente. A sobrevivência após o diagnóstico de metástases foi calculada a partir da data da confirmação histopatológica da doença metastática. Foram investigadas as associações entre o estado dos recetores no tumor primário, o estado dos recetores no tecido metastático e a conversão do estado dos recetores nas metástases com os resultados em termos de sobrevivência.

Resultados

A pleura foi o local metastático mais frequente (54,3%). Nos tumores primários, as taxas de positividade para ER, PR e HER2 foram de 87%, 63% e 21%, respetivamente, enquanto nos tecidos metastáticos foram de 69,6%, 45,7% e 25,6%. Observou-se perda de expressão de ER em 17,4% dos casos, sem se verificar qualquer ganho de expressão. A perda de expressão de PR ocorreu em 23,9% dos casos e o ganho de expressão em 6,5%, enquanto a perda e o ganho de expressão de HER2 foram observados em 7,0% e 11,6% dos casos, respetivamente. A análise univariada identificou a metastização pleural (HR = 3,442; $p=0,017$), a negatividade para ER no tecido metastático (HR = 3,306; $p=0,008$), a negatividade para PR no tecido metastático (HR = 2,793; $p=0,037$) e a perda de expressão de ER (HR = 3,095; $p=0,022$) como fatores prognósticos adversos. Na análise multivariada, a metastização pleural (HR = 4,424; $p=0,009$) e a perda de expressão de ER (HR = 3,669; $p=0,010$) mantiveram-se como preditores independentes de pior sobrevivência. Não foi encontrada qualquer associação significativa entre o estado de HER2 e a sobrevivência.

Conclusão

A perda de expressão de ER e a metastização pleural constituem fatores prognósticos adversos independentes nas metástases torácicas do cancro da mama. Estes achados, relacionados com a biologia tumoral, em particular com as alterações do estado dos recetores entre o tumor primário e as metástases, poderão contribuir para orientar abordagens terapêuticas personalizadas.

RESUMO

Introdução

O estado dos recetores pode sofrer alterações durante a progressão metastática, podendo modificar as decisões

Keywords:

Breast Neoplasms; Erb-b2 Receptor Tyrosine Kinases; Neoplasm Metastasis; Receptors, Estrogen; Receptors, Progesterone

Palavras-chave:

Metástase Neoplásica; Neoplasias da Mama; Receptor Proteína Tirocina Quinase ErbB-2; Receptores de Estrogénio; Receptores de Progesterona

INTRODUCTION

Metastatic disease is considered the leading cause of cancer-related mortality in breast cancer.¹ Accumulating evidence indicates that receptor discordance between primary and metastatic tumors may significantly affect survival outcomes. Although receptor loss has consistently been associated with adverse prognosis, receptor positivity (ER, PR, HER2) differing from that of the primary tumor may also be detected, and such differences may allow modifications in systemic therapy, potentially leading to improved survival outcomes.²⁻⁴ Accordingly, current guidelines recommend biopsy of metastatic lesions and reassessment of receptor status to optimize treatment strategies in metastatic breast cancer.^{2,5-7}

A consistent finding across studies is that receptor alterations in metastatic disease most frequently manifest as hormone receptor (ER, HR) loss, which is strongly associated with poorer survival outcomes.^{1,8,9} Equally emphasized in the literature is the clinical importance of reassessing receptor status in metastatic lesions and incorporating these findings into therapeutic decision-making.^{5,7,10}

While previous studies have investigated receptor discordance in distant metastases irrespective of organ involvement, as well as in selected organ-specific metastases (e.g., liver, bone, brain), data focusing specifically on thoracic metastases remain limited. The present study was designed to evaluate receptor profile alterations in thoracic metastases of breast cancer, to stratify patients according to receptor status and direction of conversion, to comparatively assess their impact on survival, and to identify receptor patterns associated with more favorable survival outcomes.

MATERIAL AND METHODS

In this single-center retrospective study, hospital records of 72 patients across all age groups who underwent surgical intervention for thoracic metastases of breast cancer between January 2020, and December 2024 were reviewed. Ethical approval for the study was obtained from the T.C. Istanbul Medipol University Non-Interventional Clinical Research Ethics Committee (Approval No: 976; Date: 14.08.2025). Survival analyses were conducted using follow-up data updated through July 2025. Receptor profiles and patterns of receptor conversion were categorized and analyzed to determine their association with post-metastasis survival. The impact of primary tumor receptor status, metastatic receptor status, and receptor discordance on survival outcomes was statistically assessed.

Inclusion criteria were prior diagnosis of breast cancer and surgical management of thoracic metastasis, female sex, and documented receptor evaluation in both primary and metastatic pathology specimens. Thoracic specimens included pleura, pleural

effusion, pericardium, pericardial effusion, mediastinal lymph nodes, chest wall, and lung tissue. Exclusion criteria comprised: male sex, unavailable primary tumor pathology, unavailable metastatic pathology, surgery-related in-hospital mortality, absence of ER, PR, or HER2 receptor status in pathology reports, and incomplete clinical records.

Of 72 eligible patients, one was excluded for male sex and 25 for incomplete records or unavailable primary pathology data. The final study consisted of 46 patients. HER2 status could not be determined by immunohistochemistry (IHC) in three cases, and no additional confirmatory testing was performed; therefore, HER2 analyses were conducted in 43 patients.

Statistical Analysis

All statistical analyses were performed using IBM SPSS Statistics version 25.0 (IBM Corporation, Armonk, NY, USA). Post-metastasis survival was defined as the interval between the date of histopathological confirmation of thoracic metastasis and the date of death or last follow-up. Overall survival rates, including 6-month, 1-year, and 2-year cumulative survival, as well as mean estimated



survival time, were calculated using the Kaplan–Meier method and presented with 95% confidence intervals (CIs). The effects of pleural metastasis and PR and HER2 conversion to positivity on survival were evaluated using the log-rank test. Univariate and multivariate analyses were performed using Cox proportional hazards regression models. Hazard ratios (HRs), 95% confidence intervals (CIs), and Wald statistics were calculated. Differences in receptor status distributions between primary breast tumors and corresponding metastatic tissues were assessed using the McNemar test. A p -value < 0.05 was considered statistically significant.

RESULTS

Demographic characteristics and the distribution of thoracic metastatic sites were evaluated. The mean age of the patients was 51.6 ± 13.0 years, with a range of 26–81 years. The most common sites of thoracic metastasis of breast cancer were the pleura (54.3%), followed by the lung (19.6%), lymph nodes (10.9%), pericardium (10.9%), and chest wall (4.3%) (Table 1).

Table 1 Demographic and clinical characteristics of the patients included in the study

n=46	
Age at diagnosis (years)	51.6 ± 13.0
Age range at diagnosis (years)	26–81
<i>Specimen</i>	
Pleura	25 (54.3%)
Lung	9 (19.6%)
Lymph node	5 (10.9%)
Pericardium	5 (10.9%)
Chest wall	2 (4.3%)
<i>Vital status after thoracic metastasis</i>	
Alive	26 (56.5%)
Deceased	20 (43.5%)
Follow-up duration after thoracic metastasis (months)	18 (1–67)

The median follow-up time after thoracic metastasis was 18 months (range, 1–67 months). At the end of the study period, 26 patients (56.5%) were alive, and 20 patients (43.5%) were deceased. According to Kaplan–Meier survival analysis, the 6-month cumulative survival rate was 0.761, whereas the 1- and 2-year cumulative survival rates were 0.673 and 0.539, respectively. The mean estimated overall survival was 40.2 months, with a 95% confidence interval (CI) of 31.4–49.0 months (Fig. 1).





Figure 1 Kaplan–Meier survival curve after thoracic metastasis

In the primary breast cancer tissue, ER was positive in 87% of patients, PR in 63%, and HER2 in 21%. In the thoracic metastases, ER was positive in 69.6% of patients, PR in 45.7%, and HER2 in 25.6% (Fig. 2).



^A McNemar test between primary cancer receptor positive and metastasis receptor positive

Figure 2 Distribution and changes in ER, PR, and HER2 status between primary breast tumors and corresponding metastases

Hormone receptor positivity was observed less frequently in metastatic tissue compared to primary tumors, whereas HER2 positivity was more frequent in metastatic tissue. A statistically significant difference in ER positivity was found between metastatic and primary tumor tissues ($p=0.008$). PR positivity showed a borderline difference ($p=0.057$). No significant difference was observed for HER2 status between the two groups ($p=0.727$) (Fig. 2).

Receptor Changes in Metastatic Tissue

The direction of receptor conversion from primary tumor tissue to metastatic tissue (positive to negative, negative to positive, or no change) was evaluated. ER conversion from positive to negative was observed in 17.4% of cases, whereas no negative-to-positive conversion was detected. PR conversion from positive to negative occurred in 23.9% of cases, while negative-to-positive conversion was observed in 6.5%. HER2 conversion from positive to negative was identified in 7.0% of cases, and negative-to-positive conversion in 11.6% (Fig. 2).

These findings suggest that receptor loss, particularly of hormone receptors, is common during the metastatic process. In addition, new receptor positivity may emerge in approximately 10% of cases.

Survival Analysis According to Metastatic Site

Cumulative survival was markedly lower from the early period in patients with pleural metastasis. In cases with non-pleural metastases, the 6-month, 1-year, and 2-year survival rates were 85.7%, 85.7%, and 73.5%, respectively, with a mean overall survival of 52.7 months (95% CI: 41.8–63.6). In contrast, patients with pleural metastasis had survival rates of 68.0%, 51.7%, and 37.6% at 6 months, 1 year, and 2 years, respectively, and a mean overall survival of 29.0 months (95% CI: 17.6–40.4). The difference between the groups was statistically significant (log-rank = 6.653; $p=0.010$) (Fig. 3).




Figure 3 Kaplan–Meier survival curves by metastatic site (pleural versus other)

Survival Analysis After Thoracic Metastasis

Univariate Cox Regression Analysis

Among metastatic sites, pleural involvement was identified as a significant predictor of mortality following thoracic metastasis.

Age, non-pleural metastatic sites, and primary tumor receptor status were not significantly associated with survival ($p>0.05$). Pleural metastasis was associated with a 3.4-fold increased risk of death compared to other metastatic sites (HR = 3.442, 95% CI: 1.242–9.536, $p=0.017$).

ER negativity (HR = 3.306, $p=0.008$) and PR negativity (HR = 2.793, $p=0.037$) in metastasis were significantly associated with increased mortality, whereas HER2 status was not ($p=0.988$). Loss of ER positivity during metastatic progression independently predicted worse survival (HR = 3.095, $p=0.022$). No significant association was observed for PR or HER2 conversion ($p>0.05$) (Table 2).

Table 2 Univariate Cox proportional hazards models evaluating survival after thoracic metastasis based on metastatic receptor status and receptor conversion patterns

	HR (95% CI)	Wald	p-value
Metastasis			
ER negative	3.306 (1.365–8.011)	7.014	7.014
PR negative	2.793 (1.065–7.324)	4.363	4.363
HER2 negative	0.992 (0.322–3.053)	0.001	0.001
Receptor conversion (metastatic versus primary)			
ER conversion to negative	3.306 (1.365–8.011)	5.282	5.282
PR conversion to negative	2.793 (1.065–7.324)	1.895	1.895
HER2 conversion to negative	0.992 (0.322–3.053)	0.167	0.167

HR: hazard ratio, CI: confidence interval.

Multivariate Cox Regression Analysis

In multivariate analysis, pleural metastasis independently predicted worse survival after thoracic metastasis (HR = 4.424, 95% CI: 1.461–13.395, $p=0.009$). ER negativity in metastatic tissue showed a borderline association with increased mortality (HR = 3.198, $p=0.051$).

Loss of ER positivity in the metastatic tissue independently increased mortality risk (HR = 3.669, 95% CI: 1.362–9.885, $p=0.010$). After inclusion of ER conversion in the model, pleural metastasis remained an independent predictor of death (HR = 3.238, 95% CI: 1.125–9.324, $p=0.029$) (Table 3).

Table 3 Multivariate Cox proportional hazards analysis of receptor conversion and survival after thoracic metastasis

	HR (95% CI)	Wald	p-value
Age at diagnosis	1.020 (0.981–1.062)	1.016	0.313
Pleural metastasis	3.238 (1.125–9.324)	4.743	0.029
ER conversion to negative	3.669 (1.362–9.885)	6.606	0.010

HR: hazard ratio, CI: confidence interval.

When the impact of receptor gain on survival was evaluated, no comparison could be performed for ER, as no cases demonstrated

conversion from negative to positive status. For PR, the “conversion to positive” subgroup was limited in size ($n = 3$), with 6-month, 1-year, and 2-year survival rates of 100%. However, the difference between groups was not statistically significant (log-rank = 2.062, $p=0.151$). In HER2, patients with conversion to positive status had survival rates of 60.0% at all time points, with a mean overall survival of 27.6 months (95% CI: 10.0–45.2). In contrast, the remaining group (HER2 conversion to negative status or no change) had 6-month, 1-year, and 2-year survival rates of 81.6%, 71.0%, and 58.5%, respectively, with a mean overall survival of 43.3 months (95% CI: 34.0–52.6). No statistically significant difference was observed between the groups (log-rank = 0.183, $p=0.688$).

DISCUSSION

Receptor discordance in breast cancer metastases has long been recognized, and large series published particularly after 2010s have demonstrated the association between receptor conversion and clinical outcomes, highlighting the importance of metastatic biopsy.¹¹ The primary hypothesis of our study was to prove that receptor profiles in thoracic metastases of breast cancer may differ from those of the primary tumor, with receptor conversion to positive in favor of the metastasis, potentially leading to modification of systemic therapy—the mainstay of treatment—and ultimately improving patient survival. Contrary to expectations, receptor alterations were predominantly characterized by loss of receptor expression rather than gain. Moreover, even in cases where receptor conversion resulted in newly acquired positivity, this change did not translate into a survival benefit.

In our study, no cases of ER conversion to positive status were observed; therefore, survival analysis for ER gain could not be performed. PR conversion to positive was detected in three patients, and HER2 conversion to positive in five patients. Neither PR nor HER2 gain was associated with a significant survival advantage. In contrast, the literature suggests that in cases demonstrating ER or PR gain at metastasis or recurrence, the initiation of appropriate endocrine therapy may lead to improved survival outcomes.^{12,13}

In some studies, HER2 gain in the metastatic setting has been emphasized as identifying a subgroup of patients who may benefit from anti-HER2-targeted therapy. In a pilot study including 13 patients, Francis *et al* reported HER2 conversion from negative in the primary tumor to positive in metastatic tissue in seven cases, supporting this hypothesis.¹⁴ Similarly, Niikura *et al*, by demonstrating an association between HER2 loss and poorer survival, highlighted the potential role of HER2 status changes

in guiding treatment decisions, underscoring that metastatic biopsies may serve not only for diagnostic confirmation but also for therapeutic guidance and avoidance of unnecessary toxicity and overtreatment.¹⁵ Conversely, some studies have reported no significant association between HER2 conversion and survival outcomes.^{16,17} In an organ-specific cohort focusing on breast cancer liver metastases, Sundén *et al* suggested that improved survival observed in patients with HER2 gain might be attributable to access to anti-HER2 therapies.¹⁸ In our study, neither HER2 gain nor loss was significantly associated with survival.

In a single-center retrospective cohort study published in October 2025, Yoshino *et al* evaluated 33 patients who developed lung metastases from breast cancer and underwent sampling or resection. The authors reported that anti-HER2 therapy was initiated in cases demonstrating HER2 gain, even at low expression levels. Although no statistically significant survival advantage was demonstrated, the overall interpretation of the study was that subtype conversion between primary and metastatic lesions may enable more personalized treatment strategies.¹⁹ When the association between metastatic site and survival was examined, pleural metastasis emerged as an independent risk factor in both univariate and multivariate analyses. Pleural effusion represents the most common presentation of pleural involvement in breast cancer.²⁰ As with other malignancies, pleural metastasis in breast cancer—particularly when associated with malignant pleural effusion—is generally linked to poor prognosis.^{21,22} However, despite the overall unfavorable prognostic implications of pleural metastases, Light has suggested that breast cancer effusions are associated with a more favorable survival rate when they are estrogen receptor positive and the cells show a morula clustering pattern on cytology.²³

Loss of ER expression was identified as a poor prognostic factor in our study. Consistent with our findings, the literature indicates that hormone receptor loss in metastatic disease is associated with worse survival outcomes. While some studies have evaluated ER and PR loss separately, the multicenter real-world ESME cohorts assessed hormone receptor status collectively (ER and/or PR). Grinda *et al* reported that hormone receptor loss in metastatic disease was associated with adverse survival.¹⁷ Deluche *et al* reported a median overall survival of 39.5 months and attributed improvements in 5-year survival rates compared to earlier periods primarily to advances in HER2-targeted therapies.²⁴

In a meta-analysis, Aurilio *et al* demonstrated that ER/PR loss in the metastatic setting was associated with reduced effectiveness of endocrine therapy.²⁵ Similarly, Shiino *et al* showed that patients experiencing receptor loss had significantly poorer overall survival,

suggesting that receptor loss during metastatic progression may reflect increased tumor aggressiveness and represent a key mechanism underlying resistance to endocrine therapy.³ In earlier studies, Dowling *et al* also associated receptor discordance with worse survival, attributing this effect to the absence of appropriately targeted treatment.²⁶

Limitations

The main limitations of our study are its retrospective design, single-center setting, and limited sample size. In addition, the fact that initial interventions were performed at different institutions and at different time points introduces heterogeneity in diagnostic, surgical, and histopathological techniques, thereby reducing standardization and potentially affecting data reliability. The effective sample size was reduced to approximately two thirds of the initially identified population, primarily due to missing records and limited access to complete data. Owing to incomplete documentation, analyses stratified by histological subtype could not be performed. Furthermore, the small number of cases and events in non-pleural metastatic sites limited the statistical power for subgroup analyses. The lack of information regarding stage at initial diagnosis, treatments administered after diagnosis, and the nature of surgical resection (when performed) precluded assessment of local disease control. The study evaluated overall survival (OS) only; cancer-specific survival and progression-free survival (PFS) were not assessed, preventing cause-specific mortality analyses.

CONCLUSION

Receptor profiles in thoracic metastases of breast cancer may differ from those of the primary tumor and should therefore be reassessed in the metastatic setting. Receptor conversion has the potential to influence treatment selection and may also carry prognostic implications. In particular, loss of ER expression may limit the effectiveness of endocrine therapy and necessitate consideration of alternative therapeutic strategies. The relatively frequent observation of HER2 gain is clinically relevant, as it may identify patients who could benefit from anti-HER2-directed therapies.

From a clinical perspective, pleural involvement was independently associated with a more aggressive clinical course, suggesting that this subgroup may require closer surveillance and optimized treatment strategies. Overall, our findings identify

ER loss and the presence of pleural metastasis as independent adverse prognostic factors in metastatic breast cancer with thoracic involvement. Receptor status should be re-evaluated in cases of metastatic disease. Given the potential association of pleural metastasis with poorer outcomes, early multidisciplinary evaluation, closer follow-up, and timely consideration of symptom-directed local control strategies may be warranted.

Future studies employing prospective, multicenter designs are essential to strengthen causal inference. Comprehensive data collection—including stage at initial diagnosis, menopausal status, histological subtype, immunohistochemical characteristics (receptor status, Ki-67 index), molecular subtype, treatments administered after primary diagnosis (surgery, endocrine therapy, radiotherapy, systemic therapies), degree of local disease control, and disease-free survival—would allow more robust modeling of prognostic factors. Larger prospective cohorts with sufficient event numbers are needed to clarify the potential survival impact of receptor gain. Moreover, comparative analyses of site-specific (pleura, lung, lymph node, pericardium, chest wall) receptor conversion patterns and site-specific survival outcomes in larger series would help determine any differences. The association between disease-free interval and receptor discordance remains inconclusive in the current literature and requires well-designed prospective evaluation.

Ethical Disclosures:

Conflicts of Interest: The authors have no conflicts of interest to declare.

Financing Support: This work has not received any contribution, grant or scholarship.

Confidentiality of Data: The authors declare that they have followed the protocols of their work center on the publication of patient data.

Protection of Human and Animal Subjects: The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and those of the Code of Ethics of the World Medical Association (Declaration of Helsinki as revised in 2024).

Provenance and Peer Review: Not commissioned; externally peer-reviewed.

Responsabilidades Éticas:

Conflitos de Interesse: Os autores declaram a inexistência de conflitos de interesse na realização do presente trabalho.

Fontes de Financiamento: Não existiram fontes externas de financiamento para a realização deste artigo.

Confidencialidade dos Dados: Os autores declaram ter seguido os protocolos da sua instituição acerca da publicação dos dados de doentes.

Proteção de Pessoas e Animais: Os autores declaram que os procedimentos seguidos estavam de acordo com os regulamentos estabelecidos pela Comissão de Ética responsável e de acordo com a Declaração de Helsínquia revista em 2024 e da Associação Médica Mundial.

Proveniência e Revisão por Pares: Não comissionado; revisão externa por pares.

REFERENCES

1. Yi ZB, Yu P, Zhang S, Wang WN, Han YQ, Ouyang QC, et al. Profile and outcome of receptor conversion in breast cancer metastases: a nation-wide multicenter epidemiological study. *Int J Cancer*. 2021;14:692–701. doi:10.1002/ijc.33227
2. Van Poznak C, Somerfield MR, Bast RC, Cristofanilli M, Goetz MP, Gonzalez-Angulo AM, et al. Use of biomarkers to guide decisions on systemic therapy for women with metastatic breast cancer: American Society of Clinical Oncology clinical practice guideline. *J Clin Oncol*. 2015;33:2695–704. doi: 10.1200/JCO.2015.61.1459
3. Lu Y, Tong Y, Chen X, Shen K. Association of biomarker discrepancy and treatment decision, disease outcome in recurrent/metastatic breast cancer patients. *Front Oncol*. 2021;11:638619. doi:10.3389/fonc.2021.638619
4. McAnena PF, McGuire A, Ramli A, Curran C, Malone C, McLaughlin R, et al. Breast cancer subtype discordance: impact on post-recurrence survival and potential treatment options. *BMC Cancer*. 2018;18:203. doi:10.1186/s12885-018-4101-7
5. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Breast Cancer. Version 1.2026 [accessed Jan 2026]. Available at: https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf
6. Cardoso F, Senkus E, Costa A, Papadopoulos E, Aapro M, André F, et al. 4th ESO-ESMO International Consensus Guidelines for Advanced Breast Cancer (ABC 4). *Ann Oncol*. 2018;29:1634–57. doi: 10.1093/annonc/rdy192
7. Cazzaniga ME, Pronzato P, Amoroso D, Arpino G, Atzori F, Beano A, et al. Survival outcomes of luminal metastatic breast cancer patients according to changes in molecular subtype at re-biopsy: insights from the GIM-13-AMBRA study. *Cancers*. 2025;17:1715. doi:10.3390/cancers17101715
8. Lindstrom LS, Karlsson E, Wilking UM, Johansson U, Hartman J, Lidbrink EK, et al. Clinically used breast cancer markers such as estrogen receptor, progesterone receptor, and human epidermal growth factor receptor 2 are unstable throughout tumor progression. *J Clin Oncol*. 2012; 30:2601-8. doi:10.1200/JCO.2011.37.2482
9. Shiino S, Ball G, Syed BM, Kurozumi S, Green AR, Tsuda H, et al. Prognostic significance of receptor expression discordance between primary and recurrent breast cancers: a meta-analysis. *Breast Cancer Res Treat*. 2022;191:1–14. doi:10.1007/s10549-021-06390-6
10. Duffy MJ, Harbeck N, Nap M, Molina R, Nicolini A, Senkus E, et al. Clinical use of biomarkers in breast cancer: Updated guidelines from the European Group on Tumor Markers (EGTM). *Eur J Cancer*. 2017;75:284–98. doi:10.1016/j.ejca.2017.01.017
11. Amir E, Miller N, Geddie W, Freedman O, Kassam F, Simmons C, et al. Prospective study evaluating the impact of tissue confirmation of metastatic disease in patients with breast cancer. *J Clin Oncol*. 2012;30:587-92. doi:10.1200/JCO.2010.33.5232
12. Zhao W, Sun L, Dong G, Wang X, Jia Y, Tong Z. Receptor conversion impacts outcomes of different molecular subtypes of primary breast cancer. *Ther Adv Med Oncol*. 2021;13:17588359211012982. doi:10.1177/17588359211012982
13. Shen J, Xu L, Shi J, Zhao L, Shi S, Feng J, et al. Prognostic Value and Influence of Receptor Conversion on Treatment Regimen in Metastatic Breast Cancer at the First Time of Recurrence. *Oncol Res Treat*. 2020;43:620–7. doi:10.1159/000509673
14. Francis IM, Alath P, George SS, Jaragh M, Al Jassar A, Kapila K. Metastatic breast carcinoma in pleural fluid: Correlation of receptor and HER2 status with the primary carcinoma-a pilot study. *Diagn Cytopathol*. 2016;44:980–6. doi:10.1002/dc.23607
15. Niikura N, Liu J, Hayashi N, Mittendorf EA, Gong Y,

- Palla SL, et al. Loss of human epidermal growth factor receptor 2 (HER2) expression in metastatic sites of HER2-overexpressing primary breast tumors. *J Clin Oncol*. 2012;30:593-9. doi:10.1200/JCO.2010.33.8889
16. Schrijver W, Suijkerbuijk KP, van Gils CH, van der Wall E, Moelans CB, van Diest PJ. Receptor conversion in distant breast cancer metastases: a systematic review and meta-analysis. *J Natl Cancer Inst*. 2018; 110: 568-80. doi:10.1093/jnci/djx273
17. Grinda T, Joyon N, Lusque A, Lefevre S, Arnould L, Penault-Llorca F, et al. Phenotypic discordance between primary and metastatic breast cancer in the large-scale real-life multicenter French ESME cohort. *NPJ Breast Cancer*. 2021;7:41. doi:10.1038/s41523-021-00252-6
18. Sundén M, Norgren S, Lundqvist R, Andersson A, Sund M, Hemmingsson O. Receptor conversion and survival in breast cancer liver metastases. *Breast Cancer Res*. 2023;25:105. doi:10.1186/s13058-023-01706-4
19. Yoshino R, Takahashi K, Hatanaka N, Ito A, Ujiiie N, Yasuda S, et al. Analysis of prognostic factors after pulmonary resection for metastatic breast cancer: a 23-year single-institution retrospective study. *Gen Thorac Cardiovasc Surg*. 2026;74:309-16. doi: 10.1007/s11748-025-02207-1.
20. Connolly JE Jr, Erasmus JJ, Patz EF Jr. Thoracic manifestations of breast carcinoma: metastatic disease and complications of treatment. *Clin Radiol*. 1999;54:487-94. doi:10.1016/s0009-9260(99)90844-9
21. Butow PN, Coates AS, Dunn SM. Psychosocial predictors of survival: metastatic breast cancer. *Ann Oncol*. 2000;11:469-74. Doi:10.1016/s0009-9260(99)90844-9
22. Li S, Li C, Shao W, Liu X, Sun L, Yu Z. Survival analysis and prognosis of patients with breast cancer with pleural metastasis. *Front Oncol*. 2023;13:1104246. doi.org/10.3389/fonc.2023.1104246
23. Light RW. Malignant pleural and pericardial effusions. In: Sellke FW, Rutherford RB, Brunnicardi FC, editors. *Sabiston & Spencer Surgery of the Chest*. 10th ed. Philadelphia: Elsevier; 2024. p. 554-66.
24. Deluche E, Antoine A, Bachelot T, Lardy-Cleaud A, Dieras V, Brain E, et al. Contemporary outcomes of metastatic breast cancer among 22,000 women from the multicentre ESME cohort 2008-2016. *Eur J Cancer*. 2020;129:60-70. doi :10.1016/j.ejca.2020.01.016
25. Aurilio G, Disalvatore D, Pruneri G, Bagnardi V, Viale G, Curigliano G, et al. A meta-analysis of oestrogen receptor, progesterone receptor and human epidermal growth factor receptor 2 discordance between primary breast cancer and metastases. *Eur J Cancer*. 2014;50:277-89. doi:10.1016/j.ejca.2013.10.004
26. Dowling GP, Keelan S, Cosgrove NS, Daly GR, Giblin K, Toomey S, et al. Receptor discordance in metastatic breast cancer: a review of clinical and genetic subtype alterations from primary to metastatic disease. *Breast Cancer Res Treat*. 2024;207:471-6. doi:10.1007/s10549-024-07431-6